

Effect of Foliar Application of Nitrogen and Boron on the Yield of Tomato (*Lycopersicon esculentum* Mill.)

Abhishek Tiwari¹, R.Shukla²

Department of Agricultural Sciences and Allied Industries

Rama University, Mandhana, Kanpur (U.P.)

Corresponding author: draneeta@ramauniversity.ac.in

Abstract

The present investigation entitled “Effect of Nutrient application of Nitrogen and Boron on the Yield of Tomato (*Lycopersicon esculentum* Mill.)” was conducted at the experimental farm of the faculty of Agricultural Sciences and Allied Industries, Rama University, Mandhana, Kanpur (U.P.) during 2017-18. The experiment was laid out in randomized block design (RBD) with three replications comprising of 16 treatment combinations. The treatments were Nitrogen (0 %, 0.10 %, 0.15 % and 0.20 %) and Boron (0 %, 0.15 %, 0.20 % and 0.25 %). Seeds of Azad T-9 were sown in the nursery on 15th Sep.2017 and transplanting was done on 22th Oct.,2017 and result, N3 (0.20%) and B3 (0.25%) produced maximum significant values of every yield attributes of tomato. So, it is advised to research workers and vegetable growers of Central Uttar Pradesh that for obtaining optimum yield spraying with 0.20% nitrogen and 0.25% boron is recommended.

Keywords: Spraying of fertilizer; Tomato; Nitrogen, Boron and Yield experiment

Introduction

Tomato (*Lycopersicon esculentum* Mill.) a member of Solanaceae family and is one of the most popular, nutritious and widely grown vegetable plants across the world. It is origin place in South America. Due to the excellent adaptability to wider range of soil and climatic conditions it is widely grown in any parts of the world (Ahmed, 1976). Tomato plants were brought to Europe by Red Indians and started cultivation of tomato. It was introduced in Indian Subcontinent by the Europeans. Later on, local people also started its consumption due to its popularity. It is now used everywhere in the country in so many forms. Balance fertilizations in crops will act as an insurance against possible nutrient deficiencies that may be created by the respected use of a single nutrient. Among different nutrients that were required for tomato cultivation nitrogen and boron are the most important nutrients. It also promotes vegetative growth, flower and fruit setting of tomato. Next to nitrogen fertilizer, phosphatic fertilizers dwell is the second most important input for increasing crop production.

All types of soil are suitable for tomato production including sandy and heavy clay with soil pH of 5.5 to 7.5 is best one. However, sandy loam soil is considered best for an early crop. Highest yield can be obtained by growing tomato in loam, clay loam and silt loam having enough organic matter.

Fertilizer plays important role in enhancing of tomato yield and quality. Essential macro nutrient (N, P and K) and some micro nutrient such as (B, Cu and Zn) are very important for enzymatic

reactions with in plant body such as making of RNA and DNA, protein formation, synthesis of cell wall, occurrence of flowering and fruit formation, important part of growth hormone, while their deficiency affects yield and quality of plants.

Nitrogen is essential for carbohydrates, protein metabolism and sexual fertilization. It is also essential for the synthesis of tryptophan which is the precursor of In dole Acetic Acid (IAA).

Boron is essential for yield and quality of tomato. It helps in the development of cell wall, occurrence of cell division, formation of the vascular bundle, protein synthesis, root system development, fruit and seed formation and transport of sugar.

Keeping the above points in view the present investigation entitled “Effect of Nutrient application of Nitrogen and Boron on the Yield of Tomato (*Lycopersicon esculentum* Mill.)”

Materials and methods

Site and location of experiment

The experiment was conducted in Rama university Research farm Kanpur during the year 2017-2018. Geographically Kanpur is situated in the Gangetic plains of central U.P. It lies in altitude and longitude ranges between 25.28° to 28.50° north and 79.31° to 84.34° east at elevation of 125.91 m above sea level.

Weather conditions

Kanpur is characterized by sub-tropical climate with hot dry summer and cold winters. The annual rainfall is about 800-860 mm. The major portion of rain is received between July to

September, with scattered shower in winter from the North-East monsoon.

Experiment Details

The treatment comprised combination of four levels of Nitrogen (0 %, 0.10 %, 0.15 % and 0.20 %) and Boron (0 %, 0.15 %, 0.20 % and 0.25 %). Thus, the total numbers of treatments with their symbols have been presented in Table 3.5

Table 3.5: Details of treatments with their symbols

Treatments	Symbols
N levels (%)	
0 %	N ₀
0.10%	N ₁
0.15 %	N ₂
0.20 %	N ₃
B levels (%)	
0 %	B ₀
0.15%	B ₁
0.20 %	B ₂
0.25 %	B ₃

Experiment Design and Layout

The experiment was carried out in Randomized Block Design with three replications. In all, there were sixteen treatment combinations and were randomly allotted to different plots with the help of Fisher’s random number table (Fisher, 1950). The gross plot size was 2.25 m × 1.50 m (3.375 m²). The treatment combinations are sown in Table 1

Table 1: Treatment combinations

Treatment	Symbols
N ₀ B ₀	T ₁
N ₀ B ₁	T ₂
N ₀ B ₂	T ₃
N ₀ B ₃	T ₄
N ₁ B ₀	T ₅
N ₁ B ₁	T ₆
N ₁ B ₂	T ₇
N ₁ B ₃	T ₈
N ₂ B ₀	T ₉
N ₂ B ₁	T ₁₀
N ₂ B ₂	T ₁₁
N ₂ B ₃	T ₁₂
N ₃ B ₀	T ₁₃
N ₃ B ₁	T ₁₄

N ₃ B ₂	T ₁₅
N ₃ B ₃	T ₁₆
Total number of treatments	16

Statistical Analysis

The experimental data recorded on each aspect on each treatment were statistically computed in RBD as following procedure which is given by Panse and Sukhatme (1985). For calculating standard error of mean and critical difference (t) value was taken at 0.05 level of significance.

Techniques Adopted for Recording Observations

Yield attributes-Number of fruits per plant

Several pickings were required as all the fruits did not mature at a time. In each picking, fruits were counted and after last picking, the average number of fruits per plant was calculated.

Diameter of fruit (cm):

The diameter of fruit was measured in centimetre from the middle of the fruit with the help of a Vernier Calliper. The diameter of all the fruit of sampled plants was taken and mean was worked out.

Weight of fruit (g)

The weight of randomly selected fruits was pooled and average fruit weight was calculated in gram.

Results and Discussion

1. Maximum 27.83 number of fruits per plant of tomato was observed with N₃ (0.20%) and it was minimum 23.88 with N₀. Boron at 0.25 % produced maximum 26.80 fruits per plant and it was minimum 24.71 in B₀. Interactive treatment N₃B₃ recorded maximum 29.53 fruit per plant and it was recorded minimum 21.20 with N₀B₀ as given below table-

Table 1: Analysis of variance for number of fruits per plant of tomato (*Lycopersicon esculentum* Mill.)

Source of Variation (S.V.)	Degree Of Freedom (d.f.)	Sum of square (S.S.)	Mean sum of square (M.S.S.)	F cal.	F tab.	
					At 5 %	At 1 %
Replication	2	5.230	2.615	1.454		
Factor N	3	59.598	19.861	10.962**	2.92	4.51
Factor B	3	17.445	5.819	3.230*	2.92	4.52
N × B	9	21.043	2.338	1.381	2.25	3.09
Error	30	54.523	1.815			
Total	47	157.840				

2. Diameter of fruit was obtained maximum 7.47 cm with N₃ (0.20%) and it was minimum 6.42 cm with N₀ recorded maximum 7.42 cm diameter of fruit and minimum 6.68 cm diameter of fruit showed with B₀. Interactive treatment N₃B₃ recorded maximum 7.55 cm diameter of fruits and it was minimum 5.12 with N₀B₀ as given below table-

Table: Analysis of variance diameter of fruits (cm) of tomato (*Lycopersicon esculentum* Mill.)

Source of Variation (S.V.)	Degree Of Freedom (d.f.)	Sum of square (S.S.)	Mean sum of square (M.S.S.)	F cal.	F tab.	
					At 5 %	At 1 %
Replication	2	0.610	0.310	2.899		
Factor N	3	3.223	1.133	10.666**	2.91	4.54
Factor B	3	0.977	0.356	3.045*	2.94	4.53
Zn × B	9	1.267	0.145	1.251	2.77	3.14
Error	30	3.213	0.151			
Total	47	9.466				

3. The maximum 43.99g fruit weight was obtained with N₃ (0.20%) and it was minimum 41.58g with N₀ (0.25%) recorded maximum 43.17g fruit weight and it was minimum 42.04g with B₀. Interactive treatment N₃B₃ recorded maximum 44.12g fruit weight and it was recorded minimum 40.06g in N₀B₀ as given below table-

Table -: Analysis of variance for weight of fruit (g) of tomato (*Lycopersicon esculentum* Mill.)

Source of Variation (S.V.)	Degree Of Freedom (d.f.)	Sum of square (S.S.)	Mean sum of square (M.S.S.)	F cal.	F tab.	
					At 5 %	At 1 %
Replication	2	5.656	2.828	1.326		
Factor N	3	197.261	65.736	30.814**	2.92	4.53
Factor B	3	56.936	18.971	8.892**	2.82	4.59
N × B	9	70.339	7.788	3.649*	2.23	3.11
Error	30	64.555	2.131			
Total	47	393.889				

Conclusion

On the basis of experiment conducted during Rabi season of 2017-2018 at Rama university research farm Kanpur (U.P.) with the summary of results as From above scenario of result, N₃ (0.20%) and B₃ (0.25%) produced maximum significant values of every yield attributes of tomato. So, it is advised to research workers and vegetable growers of Central Uttar Pradesh that for obtaining optimum yield spraying with 0.20% nitrogen and 0.25% boron is recommended.

References

1. Awar, R. and Karami, E. (2016). Effect of macro and micro elements foliar spray on the quality and quantity of tomato. *International*

- Journal of Agricultural Policy and Research*, vol. 4 (2) : 22-28.
2. **Bhat, F.N. and Prasad, V.M. (2004).** Studies on the effect of different levels of boron and biofertilizer on growth and yield of tomato (*Lycopersicon esculentum* Mill). *New Agriculturist*, vol. 15 (1) : 137-140.
 3. **Bibi Haleema, AbdurRab and Syed Asghar Hussain (2017)** Effect of
 4. Calcium, Boron and Zinc Foliar Application on Growth and Fruit Production
 5. of Tomato. *Sarhad Journal of Agriculture*, 34(1): 19-30.
 6. **Bhat, F.N. and Prasad, V.M. (2004).** Studies on the effect of different levels of boron and biofertilizer on growth and yield of tomato (*Lycopersicon esculentum* Mill). *New Agriculturist*, vol. 15 (1) : 137-140.
 7. **Bhatt, L. and Srivastava, B.K. (2005).** Effect of foliar application of micronutrients on physical characteristics and quality attributes of tomato (*Lycopersicon esculentum*) fruits. *Indian Journal of Agricultural Sciences*, vol. 75 (9) : 591-592.
 8. **Chandra, R.; Ram, R.B.; Prakash, J.; Nath, D.; Kumar, S. and Kumar, M. (2014).** Effect of foliar application of micro-nutrients on growth and yield components of tomato (*Lycopersicon esculentum* Mill). *Trends in Biosciences*, vol. 7 (6) : 461-464.
 9. **Das, R.C. and Patro, R.S. (1989).** Effect of micro-nutrient mixture and urea on growth, yield and quality of tomato. *Orrissa. J. of Horticulture*, vol. 17 (1-2) : 37-45.
 10. **Delibas, L. and I. Akgun. (1996).** Effects of irrigation water with high boron content on some yield characteristics of tomato. *J. Tekirdag Agril. Fac.*, 5(1/2): 135-142.
 11. **Devi, C.P.; Singh, D.K. and Jain, S.K. (2013).** Effect of foliar feeding of micronutrients on growth and yield of chilli (*Capsicum annuum* var. *accuminatum* L.) cultivar Pant C-3. *Pantnagar Journal of Research*, vol. 11 (1) : 105-111.
 12. **Dongre, S.M.; Mahorkar, V.K.; Joshi, P.S. and Deo, D.D. (2000).** Effect of micro-nutrients spray on yield and quality of chilli (*Capsicum annuum* L.) var Jayanti. *Agricultural Science Digest*, vol. 20 (2): 106-107.
 13. **Emmert, E.M. (1961).** Effect of boron, dextrose and Beta-naphthoxyacetic acid on fertilizer requirements, yield and fruit quality of tomatoes. *Proc. Amer. Soc. Horti. Sci.*, vol. 77 : 494-499.
 14. **Gjurov, S.; Gencev, S. and Gerdzikov, L. (1965).** The effect of certain microelements on the earliness and yield of glasshouse tomatoes. *Gradinarska-i-Lazaraska Nauka*, vol. 2 : 321-330.
 15. **Kaya, C.; Higgs, D. and Burtor, A. (1999).** Foliar application of iron as remedy for zinc toxic tomato plants. *J. of Plant Nutrition*, vol. 22 (12) : 1829-1887.
 16. **Mushtaq, N.; Mushtaq, F.; Khan, S.H.; Javid, R.; Rehana, M. and Bharti, V. (2016).** Effect of foliar application of Boron and Zinc on seed yield and relative economics of tomato seed (*Lycopersicon esculentum* Mill) cv. Shalimar 1. *Natural Resource Management Ecological Perspectives, Indian Ecological Society : International Conference-2016, SKUAST-Jammu (18-20 February 2016).*
 17. **Piper, C.S. (1966).** Soil and Plant Analysis. Reprint for Asia Publishing House, Bombay.
 18. **Patnaik, M.C.; Raj, G.B. and Reddy, I.P. (2001).** Response of tomato to zinc
 19. and iron. *Vegetable Science*, vol. 28 (1) : 78-79.
 20. **Sahin, S.; Gebologlu, N.; Karaman, M. R. (2015).** Interactive effect of
 21. calcium and boron on growth, quality and mineral content of tomato
 22. (*Lycopersicon esculentum* Mill). *Fresenius Environmental Bulletin*, Vol. 24 (5) : 1624-1628.
 23. **Walkley, A. and Black, I. A. (1964).** An examination of the direct method for determining soil organic matter and a proposed modification of the chromic titration method. *Soil Science*, vol. 34:29-38.
 24. **Yuanxin, L. and Junhua, W. (2009).** Effect of boron and manganese on quality and antioxidative capacity in tomato. *Acta Horticulture*, vol. 13 : 823.