

## **ANTIBACTERIAL ACTIVITY OF HONEY AGAINST CLINICAL ISOLATES OF MRSA, FROM VARIOUS CLINICAL SAMPLES AT A TERTIARY CARE CENTRE, KANPUR.**

**Deepak S<sup>1</sup>, R. Sujatha<sup>1</sup>Nashra A<sup>2</sup>**

1. Tutor, Department of Microbiology, Rama Medical College Hospital and Research Center Kanpur (India)
2. Prof and Head, Department of Microbiology, Rama Medical College Hospital and Research Center Kanpur (India)
3. Research Assistant, Central research lab, Rama Medical College Hospital and Research Center Kanpur (India)

### **ABSTRACT-**

**BACKGROUND-** The continuous use of antibiotics in clinical practice has been the direct cause of the development of multiple antibiotic resistances among bacteria causing human infection. Therefore, the need for novel alternative antimicrobial strategies has renewed interest in therapeutic use natural products of ancient remedies like turmeric, honey, ginger and others exhibiting antibacterial properties.

**Aims-** To find out the efficacy of antibacterial activity of locally available honey against Methicillin Resistant *Staphylococcus aureus* (MRSA) and Methicillin Sensitive *Staphylococcus aureus* (MSSA) isolates.

**MATERIAL AND METHODS-** A prospective study was carried out from June 2018 to December 2018 in the Department of Microbiology, RMCHRC, Mandhana Kanpur, the antibacterial activity of dabur pasteurised honey was evaluated against the bacterial strains of Methicillin Resistant *Staphylococcus aureus* and Methicillin Sensitive *Staphylococcus aureus*. Their antibacterial sensitivity pattern was tested using Kirby-Bauer disc diffusion susceptibility testing technique of CLSI along with other commonly used antimicrobials

**.RESULT:-** Out of 140 clinical samples, 59 isolates were of *Staphylococcus aureus*. This includes 22 isolates of MRSA *out of which 20(90.9%) isolates were sensitive to honey & 37 isolates of (MSSA) Methicillin sensitive S. aureus out of which 30(81.08%) isolates were sensitive to honey*. Antibiotics susceptibility testing was also carried out by using other commonly used antibiotics which include Erythromycin, Ciprofloxacin, Tetracycline, Ampicillin Levofloxacin, Clindamycin, Penicillin G, Linezolid, Vancomycin. All MRSA were sensitive to linezolid where as MSSA were sensitive to all except penicillin.

**CONCLUSION** – The excellent antibacterial activity of honey against MRSA isolates indicates the usefulness of honey against bacterial infections.

## **INTRODUCTION**

Antimicrobial agents are essentially important in reducing the global burden of infectious diseases [1]. The selection and spread of multi-resistant organisms in developing countries, which can often be traced to complex socio-economic and behavioural factors, contribute to the escalating problem of antibiotic resistance worldwide. In developing countries, the irrational use of antibiotics by health professionals, unskilled practitioners, laypersons, poor drug quality, unhygienic conditions and inadequate surveillance account for the spread of resistant bacteria. Nevertheless, misuse of antibiotics is one of the reasons for the increasing rates of resistance, especially in rural areas [2]. There is an emergence of drug resistance for the drugs like vancomycin and daptomycin leading to search for still newer drugs for combating the drug resistance in Staphylococci [3]. With the irrational and excessive use of antibiotics in underdeveloped and developing countries the developed resistance may spread in the community making the strains as super bugs causing difficulties in eradication [4]. As a result, the effectiveness of the antibiotics is diminished [5]. Therefore, the need for novel alternative antimicrobial strategies has renewed interest in natural products like turmeric, honey, ginger etc., exhibiting antibacterial properties. This situation has led to a re-evaluation of the therapeutic use of ancient remedies including honey [6-8]. Honey is well known as a magic drug for almost all kinds of diseases, not to mention the fact that many people do depend more on folk medicine and

natural remedies which are cheap that have been known for their therapeutic effects over the past decades [6].

Honey has well established function as an effective antibacterial agent with a broad spectrum of activity against Gram-positive and Gram-negative bacteria [9-11]. The application of honey can promote the healing of infected wounds that do not respond to the conventional therapy, i.e., antibiotics and antiseptics [12] including wounds infected with methicillin-resistant *S. aureus* [13,14]. Laboratory studies have revealed that honey is effective against MRSA,  $\beta$ -haemolytic streptococci and Vancomycin Resistant Enterococci (VRE) [15,16]. The beneficial role of honey is attributed to its antibacterial property with regards to its high osmolarity, acidity (low pH) and content of hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>) and non-peroxide components, i.e., the presence of phytochemical components like Methylglyoxal (MGO). The antimicrobial agents in honey are predominantly hydrogen peroxide, of which the concentration is determined by relative levels of glucose oxidase, synthesized by the bee and catalase originating from flower pollen [17].

## MATERIAL AND METHODS


A prospective study was carried out from June 2018 to December 2018 in the Department of Microbiology, RMCHRC, Mandhana Kanpur, the antibacterial activity of dabur pasteurised honey was evaluated against the bacterial strains of Methicillin Resistant *Staphylococcus aureus* and Methicillin Sensitive *Staphylococcus aureus*. Their antibacterial sensitivity pattern was tested using Kirby-Bauer disc diffusion susceptibility testing technique of CLSI along with other commonly used antimicrobials.

## RESULTS


Out of 140 clinical samples, 59 isolates were of *Staphylococcus aureus*. This includes 22 isolates of MRSA out of which 20(90.9%) isolates were sensitive to honey & 37 isolates of (MSSA) Methicillin sensitive *S. aureus* out of which 30(81.08%) isolates were sensitive to honey. Antibiotics susceptibility testing was also carried out by using other commonly used antibiotics which include Erythromycin, Ciprofloxacin, Tetracycline, Ampicillin Levofloxacin, Clindamycin, Penicillin G, Linezolid, Vancomycin. All MRSA were sensitive to linezolid where as MSSA were sensitive to all except penicillin.

TOTAL NO OF <i>Staph.aureus</i> ISOLATES	MRSA	MSSA
59	22	37
Isolates sensitive to honey	20 (90.9%)	30 (81.08%)

**Fig 1: Total Number Of Isolates**




**Fig 2:- Total isolates of Staphylococcus aureus**




**Fig 3: Staphylococcal isolates sensitive to honey**



**Fig 4:- Antimicrobial activity of honey and other common antibacterial agents on/against MRSA.**



**Fig 5: Antimicrobial activity of honey and other common antibacterial agents on/against MSSA**



**Fig 6:- Antibiotic Sensitivity pattern of Staph. aureus**

## DISCUSSION

Microbial resistance to honey has never been reported [18] which makes it a very promising topical antimicrobial agent against the infection of antibiotic-resistant bacteria (*e.g.*, MDR *S. maltophilia*) and in the treatment of chronic wound infections that do not respond to antibiotic therapy. Hence honey has been used as a last-resort medication. Manuka honey has been widely researched and its antibacterial potential is renowned worldwide. The potency of honeys, such as Tualang honey, against microorganisms suggests its potential to be used as an alternative therapeutic agent in certain medical conditions, particularly wound infection.

It has been reported that honey showed both bacteriostatic and bactericidal effect against many Gram-positive as well as Gram-negative bacteria [ 19—23] Lusby *et al* [24] reported that honeys other than the commercially available antibacterial honeys (*e.g.*, manuka honey) can have equivalent antibacterial activity against bacterial pathogens. This was similar to our study where honey possess antimicrobial property. Out of 140 clinical samples, 59 isolates were of

*Staphylococcus aureus*. This includes 22 isolates of MRSA out of which 20(90.9%) isolates were sensitive to honey & 37 isolates of (MSSA) Methicillin sensitive *S. aureus* out of which 30(81.08%) isolates were sensitive to honey. Other study also states that Honey has been reported to be effective in eradicating antibiotic resistant bacteria including MRSA [25] which is a super bug now.

Antibiotics susceptibility testing was also carried out by using other commonly used antibiotics which include Erythromycin, Ciprofloxacin, Tetracycline, Ampicillin Levofloxacin, Clindamycin, Penicillin G, Linezolid, Vancomycin. All MRSA were sensitive to linezolid where as MSSA were sensitive to all except penicillin.

It is also noted that Any zone diameter having less than 7 mm shows that the organism is resistant to the honey sample but if the zone diameter is greater than 11 mm it suggests that the microorganism is sensitive to the honey with special reference to *Pseudomonas aeruginosa* [26]. Recent reviews on the successful usage of honey as a dressing on infected wounds show that many authors support the use of honey in infected wounds and some suggest the prophylactic use of honey on the wounds of patients susceptible to MRSA and other antibiotic-resistant bacteria [27]. Well documented clinical trials and researches are going on honey and nanotechnology which may provide promising results on therapeutic use of honey in the future.

## CONCLUSION

Indeed, medicinal importance of honey has been documented in the world's oldest medical literatures, and since the ancient times, it has been known to possess antimicrobial property as well as wound-healing activity. The healing property of honey is due to the fact that it offers antibacterial activity, maintains a moist wound condition, and its high viscosity helps to provide a protective barrier to prevent infection. Its immunomodulatory property is relevant to wound repair too. The antimicrobial activity in most honeys is due to the enzymatic production of hydrogen peroxide.

It is definitely worthy to consider honey as a promising future antimicrobial to be tested and studied. Rediscovering honey as a natural remedy for wound pathogens proved its effectiveness on antimicrobial resistant strains of bacteria including MRSA. In the present study, we tried to focus more on whether honey can be used for treating Staphylococcal infections with special reference to MRSA. Honey, the nature blessed and environmental friendly product may be

elaborately used in future with some more molecular studies on its method of action as an antimicrobial agent.

## REFERENCES

1. Deb Mandal M, Mandal S. Honey: its medicinal property and antibacterial activity. *Asian Pac J Trop Biomed.* 2011;1(2):154–60.
2. Okeke IN, Lamikanra A, Edelman R. Socioeconomic and behavioral factors leading to acquired bacterial resistance to antibiotics in developing countries. *Emerging Infectious Diseases.* 1999;5(1):18-27.
3. Srinivas B, Murthy SN, Prasad U, Rao BN. Demographic profile and drug resistance pattern in methicillin sensitive staphylococcus aureus in rural and urban tertiary care centers. *Journal of Pure and Applied Microbiology.* 2015;9(4):2961-69.
4. Patel A, Chauhan BP. Antimicrobial effect of Honey on MRSA isolated from pus samples. *Int J Drug Res Tech.* 2016;6(2):58-63.
5. Levy SB, Marshall B. Antibacterial resistance worldwide: causes, challenges and responses. *Nat Med.* 2004;10:122-29.
6. Bagde AB, Sawant RS, Bingare SD, Sawai RV, Nikumbh MB. Therapeutic and nutritional values of honey [Madhu]. *Int Res J Pharm.* 2013;4(3):19-22.
7. Mandal S, Deb Mandal M, Pal NK. Synergistic anti-*Staphylococcus aureus* activity of amoxicillin in combination with *Embllica officinalis* and *Nymphae odorata* extracts. *Asian Pac J Trop Med.* 2010;3(9):711-14.
8. Mandal S, Deb Mandal M, Pal NK, Saha K. Antibacterial activity of honey against clinical isolates of *Escherichia coli*, *Pseudomonas aeruginosa* and *Salmonella enterica* serovar Typhi. *Asian Pac J Trop Med.* 2010;3(12):961-64.
9. Irish J, Blair S, Carter DA. The antibacterial activity of honey derived from Australian flora. *PLoS ONE.* 2011;6:e18229.
10. Molan PC. The antibacterial nature of honey. The nature of the antibacterial activity. *Bee World.* 1992;73(1):5–28.

11. Tan HT, Rahman RA, Gan SH, Halim AS, Hassan SA, Sulaiman SA, et al. The [1] antibacterial properties of Malaysian tualang honey against wound and enteric microorganisms in comparison to manuka honey. *BMC Complement Alternat Med.* 2009;9:34.
12. Ahmed AK, Hoekstra MJ, Hage J, Karim RB. Honey-medicated dressing: transformation of an ancient remedy into modern therapy. *Ann Plast Surg.* 2003;50:143–48.
13. Natarajan S, Williamson D, Grey J, Harding KG, Cooper RA. Healing of an MRSA-colonized hydroxyurea-induced leg ulcer with honey. *J Dermatol Treat.* 2001;12:33–36.
14. Dunford C, Cooper RA, Molan PC. Using honey as a dressing for infected skin lesions. *Nurs Times.* 2000;96:7–9.
15. Allen KL, Hutchinson G, Molan PC. Potential for using honey to treat wounds infected with MRSA and VRE. First World Healing Congress, Melbourne, Australia. 2000:10–13.
16. Kingsley A. The use of honey in the treatment of infected wound. *British J Nursing.* 2001;10:13–16.
17. Weston RJ. The contribution of catalase and other natural products to the antibacterial activity of honey: a review. *Food Chemistry.* 2000;71:235–39.
18. Dixon B. Bacteria can't resist honey. *Lancet Infect Dis.* 2003;3:116.
19. Badawy OFH, Shafii SSA, Tharwat EE, Kamal AM. Antibacterial activity of bee [20] honey and its therapeutic usefulness against *Escherichia coli* O157:H7 and *Salmonella typhimurium* infection. *Rev Sci Tech Off Int Epiz.* 2004;23(3):1011- 22.
20. Wilkinson JM. Antibacterial activity of 13 honeys against [21] *Escherichia coli* and *Pseudomonas aeruginosa*. *J Med Food.* 2005;8:100-03.
21. Ogbaje EO, Ogansanya T, Aiwerioba OIR. Conventional use of honey as [22] antibacterial agent. *Ann Afri Med.* 2006;5(2):78-81.
22. Agbagwa OE, Peterside NF. Effect of raw commercial honeys from Nigeria on [23] selected pathogenic bacteria. *Afr J Microbiol Res.* 2010;4(16):1801-03.
23. Othman AS. Antibacterial activity of bee and Yemeni Sidr honey against some [24] pathogenic bacterial species. *Int J Curr Microbiol App Sci.* 2014;3(10):1015-25.

24. Lusby PE, Coombes AL, Wilkinson JM. Bactericidal activity of different honeys against pathogenic bacteria. Arch Med Res. 2005;36:464–467
25. Nagi AA, Amghalia E, Shamsudin MN, Abdullah R, Mohammed R, Sekawi Z. Antibacterial activity of honey against methicillin resistant *Staphylococcus aureus*. Research journal of Biological Sciences. 2009;4(8):943- 47.
26. Cooper RA, Molan PC. The use of honey as an antiseptic in managing Pseudomonas infection. J Wound Care. 1999;8:161-64.
27. Molan PC. Honey as a topical antibacterial agent for treatment of infected wounds. World Wide Wounds. 2011. Available at: <http://www.worldwidewounds.com/2001/november/Molan/honey-as-topical-agent.html>. Last accessed on 18.07.2017

**CORRESPONDING AUTHOR**

Dr. R.Sujatha

Professor and Head of Department of Microbiology

Rama Medical College Hospital&amp; Research Centre, Mandhana, Kanpur, U.P.

EmailID: [drsujatha152@gmail.com](mailto:drsujatha152@gmail.com)