

Bio stimulants and Bio effect or Mediated Mitigation of A biotic Stress in Crop Plant

Kartikay Bisen

Department of Plant Pathology, Faculty of Agricultural Sciences and Allied Industries, Rama University, Kanpur-209217

Abstract

Bio stimulants are various substances which are used for promoting plant growth without being a nutrient or pesticides. These are defined as the materials, other than fertilizers, that promote plant growth when applied in low quantities. They are also classified into three major groups on the basis of their source and content. These groups include products contain hormone, humic substances and amino acid containing products. Seaweed extracts contain identifiable amounts of active plant growth substances such as auxins, cytokine's, or their derivatives'. Plant bio stimulants, also referred as agricultural bio stimulants, are group of diverse substances which can be supplemented to the close environment of plants and have positive effect on nutrient uptake and plant growth. Although, the application of plant bio stimulants in the rhizo sphere facilitates nutrient uptake and plant growth, however, many of these plant bio stimulants are reported to alleviate a biotic stresses such as salt, drought and heat stresses. Over the following years, the term 'bio stimulant' is being increasingly applied by the scientific community and literatures increasing the array of substances and of modes of actions as well. In this review,

Introduction

The term "biostimulants" was actually coined by horticulturists for denoting the various substances which are used for promoting plant growth without being a nutrient or pesticides. In a web journal named Ground Maintenance (<http://grounds-mag.com>), Zhang and Schmidt from Virginia Polytechnic Institute and State University, defined bio stimulants as "materials that, in minute quantities, promote plant growth". Kauffman et al. (2007) defined biostimulants as 'materials, other than fertilizers, that promote plant growth when applied in low quantities'. Kauffman et al. (2007) summarized bio stimulants by introducing a categorization: 'Bio stimulants in a variety of formulations are generally classified into three major groups on the basis of their source and content. These groups include products contain hormone, humic substances and amino acid containing products. Seaweed extracts contain identifiable amounts of active plant growth substances such as auxins, cytokinins, or their derivatives'. Plant bio stimulants, also referred as agricultural bio stimulants, are group of diverse substances which can be supplemented to the close environment of plants and have positive effect on nutrient uptake and plant growth (du Jardin 2012). Although, the application of plant bio stimulants in the rhizo sphere facilitates nutrient uptake and plant growth, however, many of these plant bio stimulants are reported to alleviate a biotic stresses such as salt, drought and heat stresses (Van Oosten et al., 2017). Over the following years, the term 'bio stimulant' is being increasingly applied by the scientific community and literatures increasing the

array of substances and of modes of actions as well (Calvo et al., 2014, du Jardin, 2012). Truth be told, 'bio stimulant' come into view as a flexible descriptor of any substance advantageous to plants excluding fertilizers, pesticides and soil amendments. Industries and Companies in this particular sector have formed many associations, such as 'European Bio stimulants Industry Council' (EBIC) in Europe and the 'Bio stimulant Coalition' in the USA, with primary aim to dialogue with scientists, stakeholders and regulators. Study and understanding of bio stimulants and their possible effect has been growing at a significant rate (Colla and Roupahl 2015). The effectiveness of bio stimulants, exclusively in view with plant growth promotion and nutrient uptake, has been examined (du Jardin 2015; Calvo et al. 2014). In addition to plentiful manuscripts, researchers have extensively reviewed various classes of specific bio stimulants including seaweed extracts, humic and fulvic acids, protein hydrolysates, phosphate, silicon, chitosan, PGPRs, Trichoderma, arbuscular mycorrhizal fungi.

Table 1: Various types of bio stimulants reported to alleviate a biotic stresses in crop plants

Stress	Types of Bio stimulants	Crop	References
Drought tolerance	<i>Azospirillum baselines</i>	<i>Triticum aestivum</i>	Pereyra et al. 2012; Hamaoui et al. 2001
Drought tolerance	<i>A. baselines</i>	<i>Lycopersicon lycopersicum</i>	Romero et al. 2014
Drought tolerance	<i>A. nod sum</i>	<i>Camellia sinensis</i>	Spann et al. 2011
Drought tolerance	Fulvic and humic acids	<i>Festuca arundinaceous</i>	Zhanget al. 2002; Zhang et al. 2000
Drought tolerance	Fulvic and humic acids	<i>Arachis palustris</i>	Zhang et al. 2004
Drought tolerance	Megafol	<i>L. lycopersicum</i>	Petrozza et al. 2014
Drought tolerance	Sea weed extract	<i>Spinacia oleracea</i>	Xu Cet al. 2015
Drought tolerance	Sea weed extract	<i>Spiraea nipponica</i>	Elansary et al. 2016
Drought tolerance	Sea weed extract	<i>Pittosporum eugenioides</i>	Elansary et al. 2016
Drought tolerance and ion homeostasis	Sea weed extract	<i>Vitis vinifera</i>	Mancuso et al. 2006
Drought and oxidative stress	Humic acids	<i>Oryza sativa</i>	García et al. 2012
Salt tolerance	<i>A. brasilense</i>	<i>Cicer arietinum</i>	Hamaoui et al. 2001
Salt tolerance	<i>A. brasilense</i>	<i>Vicia faba</i>	Hamaoui et al. 2001
Salt tolerance	<i>A. brasilense</i>	<i>Lactuca sativa</i>	Barassiet al. 2006; Fasciglione et al. 2015
Salt and osmotic stress	<i>A. brasilense</i>	<i>T. aestivum</i>	Fasciglione et al. 1997
Salt tolerance	<i>Pantoea dispersa/ A. brasilense</i>	<i>Capsicum annum</i>	Del Amor et al. 2012
Salt tolerance	<i>A. chrococum</i>	<i>Zea mays</i>	Rojas-Tapias et al. 2012
Salt tolerance	<i>A. chrococum</i>	<i>T. aestivum</i>	Chaudhary et al. 2013[25]
Salt tolerance	<i>A. lipoferum</i>	<i>T. aestivum</i>	Bacilio et al. 2014
Salt tolerance	<i>Hartmanni bacter diazotrophicus</i>	<i>Hordeum vulgare</i>	Suarez et al. 2015
Salt tolerance and ion homeostasis	Humic acid and phosphorous	<i>Capsicum annum</i>	Çimrin et al. 2010
Salt tolerance	Humic acids	<i>Phaseolus vulgaris</i>	Aydin et al. 2012
Salt tolerance	Protein hydrolysates	<i>Z. mays</i>	Ertani et al. 2012
Salt tolerance, cold tolerance	Protein hydrolysates	<i>L. sativa</i>	Botta et al. 2013; Lucini et al. 2015
Salt tolerance	Protein hydrolysates	<i>Diospyros kaki/D. lotus</i>	Visconti et al. 2015

Salt tolerance	<i>Rhizobium leguminosarum</i>	<i>V. faba</i>	Del Pilar Cordovilla et al. 1999
Salt tolerance	<i>R. leguminosarum</i>	<i>Pisumsativum</i>	Del Pilar Cordovilla et al. 1999
Salt tolerance	Sea weed extract	<i>Poapratensis</i>	Nabati et al. 1994
Cold tolerance	<i>A. nodosum</i>	<i>Kappaphycusalvarezii</i>	Loureiro et al. 2014
Cold tolerance	<i>Burkholderia phytofirman</i>	<i>Vitisvinifera</i>	Fernandez et al. 2012; Theocharis et al. 2012
Cold tolerance	<i>Flavobacterium glaciei</i>	<i>Solanum lycopersicum</i>	Subramanian et al. 2016[34]
Cold tolerance	<i>Pseudomonas frederiksbergensis</i>	<i>Solanum lycopersicum</i>	Subramanian et al. 2016
Cold tolerance	<i>P. vancouverensis</i>	<i>Solanum lycopersicum</i>	Subramanian et al. 2016
Cold tolerance	<i>P. dispersa</i>	<i>T. aestivum</i>	Selvakumar et al. 2008
Cold tolerance	Sea weed extract	<i>Z. mays</i>	Bradáčová et al. 2016
Cold tolerance	Sea weed extract	<i>Arabidopsis thaliana</i>	Nair et al. 2012; Rayirath et al. 2009
Heat tolerance	<i>P. putida</i>	<i>T. aestivum</i>	Ali et al. 2011
Heat tolerance	<i>P. putida</i>	<i>Sorghum bicolor</i>	Ali et al. 2009
Heat tolerance	Protein hydrolysates	<i>Loliumperenne</i>	Botta et al. 2013
Heat tolerance	Sea weed extract	<i>Agrostis stolonifera</i>	Zhang et al. 2008
Ion homeostasis	<i>A. nodosum</i>	<i>Prunusdulcis</i>	Saa et al. 2015
Ion homeostasis	SWE	<i>L. sativa</i>	Möller et al. 1998
Ion homeostasis	Protein hydrolysates	<i>H. vulgare</i>	Cuin et al. 2007
Temperature tolerance	<i>A. chroococcum</i>	<i>T. aestivum</i>	Egamberdiyeva et al. 2009; Egamberdiyeva et al. 2004
Chilling stress	Glycinebetaine	<i>L. lycopersicum</i>	Park et al. 2004
Chilling tolerance	Melatonin	<i>Z. mays</i>	Kołodziejczyk et al. 2016
Heavy metal tolerance	Protein hydrolysates	<i>T. aestivum</i>	Zhu et al. 2006

Algal Extract

Marine algae commonly known as seaweeds constitute a vital part of marine coastal ecosystem. These seaweeds comprise of the multicellular marine algae that frequently reside in the coastal area of the oceans where proper substrata present. Approximately about 9,000 species of macroalgae have been estimated and are broadly classified into three primary categories based on their pigments. These classes are brown, red, and green algae or Phaeophyta, Rhodophyta,

and Chlorophyta; respectively. Among them second most abundantly present group having approximately 2000 species is brown seaweeds. These seaweeds attain their utmost biomass levels near rocky shores of the temperate zones. They are the most commonly used seaweeds in agriculture (Blunden and Gordon 1986) and, among them, *Ascophyllum nodosum* (L.) Le Jolis is the most investigated species and used in agriculture (Ugarte et al. 2012).

In addition to *A. nodosum*, other species of brown algae such as *Laminaria* spp., *Fucus* spp.,

Sargassum spp. and *Turbinaria* spp. are currently being used as bio fertilizer in agriculture (Hong et al. 2007). These seaweeds have a potential of prominent source of organic matter and nutrient and thus have been used as soil conditioner for the centuries (Mettinget al. 1988; Temple and Bomke 1988). Approximately 15 million MT of seaweeds are produced annually, and a significant portion of these products are used as bio stimulants and bio fertilizers. A range of commercially available SWEs are being used in agricultural and horticultural plants (Table 2). Extensive researches and studies have revealed the various beneficial effects of SWEs on plant health and growth, improved crop yield and elevated tolerance to the biotic and abiotic stresses (Beckett and van Staden 1989; (Fig. 1).

SWEs showed enhanced cotyledon growth resemblance to potassium fertilization (Mollar et al. 1998) Grape plants subjected to the foliar application with marine bioactive iso propanol extract of microalgae showed increased stomatal conductance and leaf water potential under drought stress.




Figure 1: Main key mechanism by seaweed extracts

The role of SWEs in alleviating the cold stress in crop plants has been studied. In recent investigations the research work has focused on SWEs and their potential role in enhancing tolerance to the chilling stress. Bradáčová et al. 2013, has tested multiple extracts for their capacity to enhance cold tolerance in maize. He reported that the extract rich in Zn and Mn were only able to mitigate the cold stress via enhanced ROS response. These outcomes specify that nutrient deficiency in plant induced by cold stress can be overcome by applying SWEs to advance oxidative stress tolerance. In prior research with corn seedlings subjected to root chilling stress and treated with essential micronutrients confirmed the efficacy of nutrient seed priming (Imran et al. 2013)

SWEs have been extensively used in horticultural crops and trees. *A. nodosum* extract is reported to increase the fresh and dry weight in spinach plant under drought stress. Lettuce plant treated with

Table 2: Commercially available seaweed products used in agriculture

Seaweed Name	Product Name	Manufacturing Company	Application
<i>Ascophyllum nodosum</i>	Acadian®	Acadian Agritech	Plant Growth stimulant
<i>A. nodosum</i>	Agri-Gro Ultra	Agri-Gro Marketing Inc.	Plant Growth stimulant
<i>Lithothamnioncalcareum</i>	Acid Buf	Chance & Hunt Ltd.	Plant Growth stimulant
<i>A. nodosum</i>	Espoma	The Espoma Company	Plant Growth stimulant
<i>A. nodosum</i>	Alga-A-Mic	Biobizz Worldwide N.V.	Plant Growth stimulant
<i>A. nodosum</i>	Kelpro	TecniprosesosBiologicos, S.A. de C.V.	Plant Growth stimulant
<i>A. nodosum</i>	Kelprosoil	Productos del Pacifico S.A. de C.V.	Plant Growth stimulant
<i>A. nodosum</i>	Stimplex	Acadian Agritech	Plant Growth stimulant
<i>A. nodosum</i>	Synergy	Green Air Products Inc.	Plant Growth stimulant
<i>A. nodosum</i>	Tasco	Acadian Agritech	Plant Growth stimulant
<i>A. nodosum</i>	Maxicrop	Maxicrop USA Inc.	Plant Growth stimulant
<i>A. nodosum</i>	Nitrozime	Hydrodynamics International Inc.	Plant Growth stimulant
<i>Microcystispyrifera</i>	Agro-Kelp	Algas y BioderivadosMarinos, S.A. de C.V.	Plant Growth stimulant
Red Marine Algae	Emerald RMA	Dolphin Sea Vegetable Company	Plant Growth stimulant
<i>Durvilleapotatorum</i>	Seasol	Season International Pty Ltd.	Plant Growth stimulant
<i>Ecklonia maxima</i>	Kelpak	BASF	Plant Growth stimulant
<i>Durvilleaantarctica</i>	Profert	BASF	Plant Bio stimulant

Conclusion

Abusive use of synthetic chemicals in agriculture has led to the hazardous effect on the human, animal and ecosystem health. Safeguarding the food security for increasing population while maintaining the ecosystem harmony is the prime challenge for the scientists. Crop production is hampered by various biotic and abiotic stresses. Various strategies have been used to alleviate the biotic stresses in crop plants. Various bio stimulants, mainly seaweed extracts have been reported to mitigate the biotic stresses in crops. These SWEs have provided the sustainable solution to the crop management under a biotic stresses. Further investigation and studies is required to understand the mechanisms of these SWEs and other bio stimulants.

References

1. Ali SKZ, Sandhya V, Grover M, Kishore N, Rao LV, Venkateswarlu B. *Pseudomonas* sp. strain AKM-P6 enhances tolerance of sorghum seedlings to elevated temperatures. *Biol Fertil Soils*. 2009; 46:45–55.
2. Ali SKZ, Sandhya V, Grover M, Linga VR, Bandi V. Effect of inoculation with a thermotolerant plant growth promoting *Pseudomonas putida* strain AKMP7 on growth of wheat (*Triticum* spp.) under heat stress. *J Plant Interact*. 2011; 6(4):239–46.
3. Ayden A, Kant C, Turin M. Humic acid application alleviates salinity stress of bean (*Phaseolus vulgaris* L.) plants decreasing membrane leakage. *Afr J Agric Res*. 2012; 7(7):1073–86.
4. Bacilio M, Rodriguez H, Moreno M, Hernandez J-P, Bashan Y. Mitigation of salt stress in wheat seedlings by a gfp-tagged *Azospirillum lipoferum*. *Biol Fertil Soils*. 2004;40(3):188–93

5. Barassi CA, Ayrault G, Creus CM, Sueldo RJ, Sobrero MT. Seed inoculation with *Azospirillum* mitigates NaCl effects on lettuce. *SciHortic*. 2006; 109(1):8–14.
6. Beckett RP, Van Staden J. The effect of seaweed concentrate on the growth and yield of potassium stressed wheat. *Plant and soil*. 1989 May 1;116(1):29-36.
7. Botta A. Enhancing plant tolerance to temperature stress with amino acids: an approach to their mode of action. *Acta Horticulturae*.2013. doi:10.17660/ActaHortic.2013.1009.1.
8. Bradáčová K, Weber NF, Morad-Talab N, Asim M, Imran M, Weinmann M, et al. Micronutrients (Zn/Mn), seaweed extracts, and plant growth-promoting bacteria as cold-stress protectants in maize. *Chem Biol Technol Agric*. 2016; 3(1):19.
9. Calvo P, Nelson L, Klopper JW. Agricultural uses of plant biostimulants. *Plant and soil*. 2014 Oct 1;383(1-2):3-41.
10. Chaudhary D, Narula N, Sindhu SS, Behl RK. Plant growth stimulation of wheat (*Triticum aestivum* L.) by inoculation of salinity tolerant *Azotobacter* strains. *Physiol Mol Biol Plants*. 2013;19(4):515–9
11. Çimrin KM, Türkmen Ö, Turan M, Tuncer B. Phosphorus and humic acid application alleviate salinity stress of pepper seedling. *Afr J Biotechnol [Internet]*. 2010; 9(36). <http://www.ajol.info/index.php/ajb/article/view/92903>. Accessed 16 Nov 2015.
12. Colla G, Rouphael Y. Biostimulants in horticulture. *Scientia Horticulturae*. 2015; 196:1-34.
13. Creus CM, Sueldo RJ, Barassi CA. Shoot growth and water status in *Azospirillum*-inoculated wheat seedlings grown under osmotic and salt stresses. *Plant PhysiolBiochem*. 1997; 35(12):939–44.
14. Cuin TA, Shabala S. Amino acids regulates salinity-induced potassium efflux in barley root epidermis. *Planta*. 2007; 225(3):753–61.
15. Del Amor FM, Cuadra-Crespo P. Plant growth-promoting bacteria as a tool to improve salinity tolerance in sweet pepper. *Funct Plant Biol*. 2012; 39(1):82–90.
16. Del Pilar Cordovilla M, Berrido SI, Ligeró F, Lluch C. Rhizobium strain effects on the growth and nitrogen assimilation in *Pisumsativum* and *Viciafaba* plant growth under salt stress. *J Plant Physiol*. 1999; 154(1):127–31.
17. Du Jardin P. Plant biostimulants: definition, concept, main categories and regulation. *Scientia Horticulturae*. 2015 Nov 30;196:3-14.
18. DuJardin P. The Science of Plant Bio stimulants—A bibliographic analysis, Ad hoc study report. European Commission; 2012.
19. Egamberdiyeva D, Höflich G. Effect of plant growth-promoting bacteria on growth and nutrient uptake of cotton and pea in a semi-arid region of Uzbekistan. *J Arid Environ*. 2004;56(2):293–301
20. Egamberdiyeva D, Höflich G. Influence of growth-promoting bacteria on the growth of wheat in different soils and temperatures. *Soil BiolBiochem*. 2003; 35(7):973–8.
21. Elansary HO, Skalicka-Woźniak K, King IW. Enhancing stress growth traits as well as photochemical and antioxidant contents of *Spiraea* and *Pittosporum* under seaweed extract treatments. *Plant Physiol Biochem*. 2016;105:310–20
22. Fasciglione G, Casanovas EM, Quillehauquy V, Yommi AK, Goñi MG, Roura SI, et al. *Azospirillum* inoculation effects on growth, product quality and storage life of lettuce plants grown under salt stress. *Sci Hortic*. 2015; 195:154–62.
23. Fernandez O, Theocharis A, Bordiec S, Feil R, Jacquens L, Clément C, et al. *Burkholderia phytofirmans* PsJN acclimates grapevine to cold by modulating carbohydrate metabolism. *MPMI*. 2012; 25(4):496–504.
24. García AC, Santos LA, Izquierdo FG, Sperandio MVL, Castro RN, Berbara RLL. Vermicomposthumic acids as an ecological pathway to protect rice plant against oxidative stress. *Ecol Eng*. 2012;47:203–8
25. Hamaoui B, Abbadi J, Burdman S, Rashid A, Sarig S, Okon Y. Effects of inoculation with *Azospirillum* brasilense on chickpeas (*Cicer arietinum*) and faba beans (*Vicia faba*) under different growth conditions. *Agronomie*. 2001;21(6–7):553–60
26. Hong DD, Hien HM, Son PN. Seaweeds from Vietnam used for functional food, medicine and biofertilizer. *Journal of Applied Phycology*. 2007 Dec 1; 19(6):817-26.
27. Imran M, Mahmood A, Römheld V, Neumann G. Nutrient seed priming improves seedling development of maize exposed to low root zone temperatures during early growth. *Eur J Agron*. 2013; 49:141–8.
28. Kauffman GL, Kneivel DP, Watschke TL. Effects of a biostimulant on the heat tolerance associated with photosynthetic capacity, membrane thermostability, and polyphenol production of perennial ryegrass. *Crop Sci* 2007 Jan; 47(1):261-7.
29. Kołodziejczyk I, Dzitko K, Szewczyk R, Posmyk MM. Exogenous melatonin improves corn (*Zea mays* L.) embryo proteome in seeds subjected to chilling stress. *J Plant Physiol*. 2016;1(193):47–56.
30. Loureiro RR, Reis RP, Marroig RG. Effect of the commercial extract of the brown alga *Ascophyllum nodosum* Mont. on *Kappaphycus salvarezii* (Doty) Doty ex P.C. Silva in situ submitted to lethal temperatures. *J Appl Phycol*. 2014;26(1):629–34.
31. Lucini L, Rouphael Y, Cardarelli M, Canaguier R, Kumar P, Colla G. The effect of a plant-derived biostimulant on metabolic profiling and crop performance of lettuce grown under saline conditions. *SciHortic*. 2015; 23(182):124–33.
32. Mancuso S, Azzarello E, Mugnai S, Briand X. Marine bioactive substances (IPA extract) improve foliar ion uptake and water stress tolerance in potted *Vitisvinifera* plants. *AdvHortic Sci*. 2006; 20(2):156–61.
33. Metting B, Rayburn WR, Reynaud PA. Algae and agriculture. Algae and human affairs. Cambridge University Press, Cambridge, UK. 1988:335-70.
34. Möller M, Smith ML. The significance of the mineral component of seaweed suspensions on lettuce (*Lactucasativa* L.) seedling growth. *J Plant Physiol*. 1998; 153(5–6):658–63.
35. Nabati DA, Schmidt RE, Parrish DJ. Alleviation of salinity stress in Kentucky bluegrass by plant growth regulators and iron. *Crop Sci*. 1994; 34(1):198–202.

36. Nair P, Kandasamy S, Zhang J, Ji X, Kirby C, Benkel B, et al. Transcriptional and metabolomic analysis of *Ascophyllumnodosum* mediated freezing tolerance in *Arabidopsis thaliana*. BMC Genom. 2012; 13(1):643.
37. Park E-J, Jeknić Z, Sakamoto A, DeNoma J, Yuwansiri R, Murata N, and et al. Genetic engineering of glycinebetaine synthesis in tomato protects seeds, plants, and flowers from chilling damage. Plant J. 2004;40(4):474–87.
38. Pereira MA, García P, Colabelli MN, Barassi CA, Creus CM. A better water status in wheat seedlings induced by *Azospirillum* under osmotic stress is related to morphological changes in xylem vessels of the coleoptile. Appl Soil Ecol. 2012;53(1):94–7.
39. Petrozza A, Santaniello A, Summerer S, Di Tommaso G, Di Tommaso D, Paparelli E, et al. Physiological responses to Megafol® treatments in tomato plants under drought stress: a phenomic and molecular approach. SciHortic. 2014;22(174):185–92
40. Rayirath P, Benkel B, Mark Hodges D, Allan-Wojtas P, MacKinnon S, Critchley AT, et al. Lipophilic components of the brown seaweed, *Ascophyllumnodosum*, enhance freezing tolerance in *Arabidopsis thaliana*. Planta. 2009;230(1):135–47.
41. Rojas-Tapias D, Moreno-Galván A, Pardo-Díaz S, Obando M, Rivera D, Bonilla R. Effect of inoculation with plant growth-promoting bacteria (PGPB) on amelioration of saline stress in maize (*Zea mays*). Appl Soil Ecol. 2012; 61:264–72.
42. Romero AM, Vega D, Correa OS. *Azospirillumbrasileense* mitigates water stress imposed by a vascular disease by increasing xylem vessel area and stem hydraulic conductivity in tomato. Appl Soil Ecol. 2014;82:38–43.
43. Saa S, Olivos-Del Rio A, Castro S, Brown PH. Foliar application of microbial and plant based biostimulants increases growth and potassium uptake in almond (*Prunusdulcis* [Mill.] D. A. Webb). Front. Plant Sci. 2015; 6:87.
44. Selvakumar G, Kundu S, Joshi P, Nazim S, Gupta AD, Mishra PK, et al. Characterization of a cold-tolerant plant growth-promoting bacterium *Pantoeadispersa* 1A isolated from a sub-alpine soil in the North Western Indian Himalayas. World J Microbiol Biotechnol. 2008; 24(7):955–60.
45. Spann TM, Little HA. Applications of a commercial extract of the brown seaweed *Ascophyllumnodosum* increases drought tolerance in container-grown “hamlin” sweet orange nursery trees. Hort Science. 2011; 46(4):577–82.
46. Suarez C, Cardinale M, Ratering S, Steffens D, Jung S, Montoya AMZ, et al. Plant growth-promoting effects of *Hartmannibacterdiazotrophicus* on summer barley (*Hordeum vulgare* L.) under salt stress. Appl Soil Ecol. 2015; 95:23–30.