Effect of nitrogen level on growth and yield Attributing characters of radish (*Raphanussativus*L.) cv. PusaChetki

Vinay J Silas, Jitendra Kumar, Sarvesh Kumar & Syed MohdQuatadah

Assistant Professor (Rama University, Mandhana ,Kanpur)

Co-responding authors: vinay.fas@ramauniversity.ac.in

Abstract

The present investigation entitled ***"Effect of nitrogen level on growth and yield Attributing character of radish (RaphanussativusL.) cv. PusaChetki "****was carried out during Rabi season in the year 2023-24 at Research plot of Department of Vegetable Science at Rama University Kanpur. The experiment was laid out in Randomised Block Design with eight treatments randomized thrice. PusaChetki variety was taken to study best suitable combination of Nitrogen to get highest yield in radish. The Different Nitrogen level used viz., N0 (Control); N1 (60 kg Nitrogen/ha); N2 (80 kg Nitrogen/ha); N3 (100 kg Nitrogen/ha); N4 (120 kg Nitrogen/ha); N5 (140 kg Nitrogen/ha); N6 (160 kg Nitrogen/ha) and N7 (180 kg Nitrogen/ha).It is observed from the investigation that the treatment N4- 120 kg Nitrogen/ha gave the highest growth and yield parameters viz., Plant height at 15 DAS, 30 DAS and at harvest(13.84,26.50 and 65.10cm), number of leaves per plant at 15 DAS, 30 DAS and at harvest(6.72,9.84and 17.10), Fresh weight of leaves at 15 DAS, 30 DAS and at harvest (44.10,83.28 and 118.32gm), Leaf area per plant at 15DAS, 30 DAS and at harvest (41.20,132.54 and 170.64cm2), Length of leaves at harvesting(38.22cm), Root length (26.78cm), Root girth (12.10cm), Average plant weight at Harvesting (224.15gm), Fresh root weight (150.45gm/plant), Yield per plot (34.15kg) and Yield per hectare (376.52q) respectively. While, the minimum results were found under N0 (Control) for all the parameters. So, we can suggest to farmers that the dose of 120 kg Nitrogen/ha gave the better results in term of growth and yield of Radish.*

Keywords: Radish cv. PusaChetki, Nitrogen, Growth and Yield parameters.

Introduction:

Radish (Raphanussativus L.) is a widely recognized root vegetable cultivated as an annual or biennial plant within the Brassicaceae family and Raphanus genus. Its appeal stems from its adaptable nature to various climates, straightforward cultivation techniques, and versatile uses, primarily focusing on its tender taproot. Both the root and leaves of radish possess low caloric value. Nitrogen stands out as a crucial element for plant growth, being integral to numerous plant structures and essential for internal and external metabolic functions. Healthy plants typically maintain nitrogen levels ranging from 3-4% in their above-ground tissues, a concentration notably higher than other nutrients. Nitrogen plays a pivotal role among macronutrients in facilitating the growth and development of plants, forming key components of vital compounds such as proteins, nucleic acids, chlorophyll, and enzymes. Soil deficient in nitrogen leads to diminished crop harvests in terms of size, weight, and quality. To augment crop production and optimize nitrogen utilization on a global scale, nitrogen fertilization remains a widely adopted practice. In radish cultivation, nitrogen fertilization holds particular significance, with its application tailored not only to maximize yield size but primarily to enhance overall yield quality. Although nitrogen proves indispensable as a constituent of essential plant components like proteins, nucleic acids, chlorophyll, and enzymes, an imbalance in its application can lead to adverse effects. Insufficient nitrogen results in poor crop yields of inferior quality, while excessive usage negatively impacts both crop quality and yield. Specifically for leafy vegetable production such as radish, nitrogen application is crucial for sustaining robust

growth, maximizing yield, and ensuring high-quality produce.

The PusaChetki variety of radish was bred at the Indian Agricultural Research Institute (IARI) in New Delhi. It yields medium-long, white roots with a desirable texture. Notably, this variety exhibits resilience to high temperatures, making it suitable for planting from mid-March to mid-August in the northern and central regions of India. In the southern and western parts of the country, it can be cultivated year-round. PusaChetki demonstrates effective seed setting in the plains, particularly due to its tendency to bolt early during October-November. Under optimal conditions, it can yield approximately 250-500 quintals of fresh radish per hectare.

Materials and Methods

LocationofExperimentalsite:

TheexperimentwascarriedoutatAgriculture ResearchFarmofRamaUniversity, Mandhana, Kanpur, India, during 2022-2023 Geographically, Kanpurdistrict falls in semiarid subtropical zone and is situated between 25.26" and 26.58"north latitude and 79.31" and 80.34" east longitude. It is located at an elevation of125.9"metersabovemeansealevel.

Seed rate and sowing:

The pure, healthy, disease and insect free vigorous and good quality seed of PusaChetki variety was used for sowing (@10 kg ha).The sowing process involved manual dibbling, ensuring a row spacing of 35 centimeters and a plant-to-plant distance of 10 centimeters. Seeds were sown at a depth of 1.25 centimeters to promote optimal root development, followed promptly by irrigation to facilitate germination.

Nutrient Management:

The only sources of nitrogen, phosphorus, and potassium utilized were urea, triple superphosphate (applied at 125 kg per hectare), and muriate of potash (applied at 175 kg per hectare), respectively. Additionally, well-decomposed cow dung was applied at a rate of 12 tonnes per hectare to the field. During land preparation, the total amount of cow dung and half of the total doses of urea and muriate of potash, along with triple superphosphate, were incorporated. The remaining portions of urea and muriate of potash were applied after 35 days from the date of seed sowing. Intercultural activities such as thinning, weeding, irrigation, and pest management were carried out as needed to optimize crop growth.

Treatmentcombinations:

Treatments Symbolused	Treatmentsdetails
N_0	Control
N_1	60 Kg/ha
N_2	80 Kg/ha
N_3	100 Kg/ha
N_4	120 Kg/ha
N_5	140 Kg/ha
N_6	160 Kg/ha
N_7	180 Kg/ha

Results and Discussion

Growth Parameters:

In the present study, the maximum Plant height (13.84 cm) at 15 DAS was recorded with the treatment **N_4**- 120 kg Nitrogen/ha. While, the minimum plant height (7.20 cm) at15 DAS was recorded with the treatment **N_0**- Control. The maximum Plant height (26.50 cm) at 30 DAS was recorded with the treatment **N_4**- 120 kg Nitrogen/ha. However, the minimum plant height (16.38 cm) at 30 DAS was recorded with the treatment **N_0**- Control. The maximum Plant height (65.10 cm) at harvest was recorded with the treatment **N_4**- 120 kg Nitrogen/ha. Moreover, the minimum plant height (35.30 cm) at harvest was recorded with the treatment **N_0**- Control. This clearly supports the fact that higher application of nitrogen has a vital role in betterment of plant physiological process such as cell division, cell elongation along with timely metabolic processes and also favored the greater assimilation of photosynthates. Similar result reported by **Moniruzzaman*et al.* (2013), Tripathi*et al.* (2017), Baloch *et al.* (2014), Pathak *et al.* (2017) and Dash *et al.* (2018).**

Similarly,The maximum numberofleavesperplant(6.72) at 15 DAS was recorded with the treatment **N_4**- 120 kg Nitrogen/ha. While, the minimum numberofleavesperplant(4.38) at15 DAS was recorded with the treatment **N_0**- Control. The maximum numberofleavesperplant(9.84) at 30 DAS was recorded with the treatment **N_4**- 120 kg Nitrogen/ha. However, the minimum numberofleavesperplant(6.16) at 30 DAS was recorded with the treatment **N_0**- Control. The maximum numberofleavesperplant(17.10) at harvest was recorded with the treatment **N_4**- 120 kg Nitrogen/ha. Moreover, the minimum numberofleavesperplant(8.96) at harvest was recorded with the treatment **N_0**- Control. This could be attributed to the enhanced availability of nutrients at the appropriate time, which might have resulted in increased photosynthetic rate and accumulation of metabolites in plants. Similarly type of findings were also reported by **Moniruzzaman*et al.* (2013), Tripathi*et al.* (2017), Thapa*et al.*(2013) and Yuan *et al.(*2015).**

The maximum Fresh weight of leaves (44.10 gm) at 15 DAS was recorded with the treatment **N_4**- 120 kg Nitrogen/ha followed. While, the minimum Fresh weight of leaves (26.25 gm) at15 DAS was recorded with the treatment **N_0**- Control. The maximum Fresh weight of leaves (83.28 gm) at 30 DAS was recorded with the treatment **N_4**- 120 kg Nitrogen/ha. However, the minimum Fresh weight of leaves (37.76 gm) at 30 DAS was recorded with the treatment **N_0**- Control. The maximum Fresh weight of leaves (118.32 gm) at harvest was recorded with the treatment **N_4**- 120 kg Nitrogen/ha. Moreover, the minimum Fresh weight of leaves (60.21 gm) at harvest was recorded with the treatment **N_0**- Control. The increase in fresh weight of leaves of the plant could be due to higher uptake and accumulation of greater amount of photosynthates. Similar results were observed by **Moniruzzaman*et al.* (2013),Jilani*et al.* (2010),Sumagaysay(2014) and Krishnakant*et al.* (2018)**.

The maximum Leaf area per plant (41.20 cm^2) at 15 DAS was recorded with the treatment **N_4**- 120 kg Nitrogen/ha. While, the minimum Leaf area per plant (30.15cm^2) at15 DAS was recorded with the treatment **N_0**- Control followed. The maximum Leaf area per plant (132.54 cm^2) at 30 DAS was recorded with the treatment **N_4**- 120 kg Nitrogen/ha. However, the minimum Leaf area per plant (65.86 cm^2) at 30 DAS was recorded with the treatment **N_0**- Control. The maximum Leaf area per plant (170.64 cm^2) at harvest was recorded with the treatment **N_4**- 120 kg Nitrogen/ha. Moreover, the minimum Leaf area per plant (78.36 cm^2) at harvest was recorded with the treatment **N_0**- Control.

Increased leaf area could be attributed to the better availability of nutrients for growth and development of plant, which, in turn, enhances the photosynthesis of leaves. The present finding also corroborate the findings of **Thakar*et al.* (2006), Sharma *et al.* (2013), Pathak *et al*. (2017), Srinivas and Naik (1990)**.

The maximum Length of leaves at harvesting (38.22 cm) was recorded with the treatment **N_4**- 120 kg Nitrogen/ha. While, the minimum Length of leaves at harvesting (22.36 cm) was recorded with the treatment **N_0**- Control. The increase in leaf length may be due to increased availability of higher dose of nutrients. Similar results were observed by **Tripathi*et al.* (2017),Jilani*et al.* (2010),Sumagaysay (2014) and Krishnakant*et al.* (2018)**.

Yield Parameters:

In the present investigation, the maximum Root length (26.78 cm) was recorded with the treatment **N_4**- 120 kg Nitrogen/ha. However, the minimum Root length (13.94 cm) was recorded with the treatment **N_0**- Control. The maximum Root girth (12.10 cm) at harvest was recorded with the treatment **N_4**- 120 kg Nitrogen/ha. Moreover, the minimum Root girth (5.21 cm) was recorded with the treatment **N_0**- Control. The root length and root girth were varied significantly among different nutrition levels. The increased Nitrogen levels produced good result in increasing the root length and root girth. Similar findings were also observed by **Moniruzzaman*et al.* (2013), Tripathi*et al.* (2017),Jilani*et al.* (2010),Thapa*et al.* (2003) and Thakar*et al.* (2006)**.

Similarly, the maximum Average plant weight at Harvesting (224.15 gm) was recorded with the treatment **N_4**- 120 kg Nitrogen/ha. While, the minimum Average plant weight at Harvesting (124.42 gm) was recorded with the treatment **N_0**- Control. The increase in Average plant weight at Harvesting may be due to increased availability of higher dose of nutrients. The maximum Fresh root weight (150.45 gm/plant) was recorded with the treatment **N_4**- 120 kg Nitrogen/ha. However, the minimum Fresh root weight (105.14 gm/plant) was recorded with the treatment **N_0**- Control. The maximum Yield per plot (34.15 kg) was recorded with the treatment **N_4**- 120 kg Nitrogen/ha. However, the minimum Yield per plot (15.26 kg) was recorded with the treatment **N_0**- Control. The maximum Yield per hectare (376.52 q) was recorded with the treatment **N_4**- 120 kg Nitrogen/ha. However, the minimum Yield per hectare (168.24 q) was recorded with the treatment **N_0**- Control. Root weight of radish increased with increased dose of nitrogen. But, after a limit the response of nitrogen was not observed. This indicates that excess application of nitrogen in radish may increase the root length but did not increase the root weight and total plant weight. This result indicate indiscriminate use of very higher doses of nitrogen did not increase root yield rather increases mere vegetative growth as compared to increase in weight of radish root. This increase in yield as well as yield attributes by different Nitrogen level may be due to increase vegetative growth giving better opportunities for photosynthetic activities and consequently increasing carbohydrates in the root resulting high yield. Similarly type of findings were also reported by

Moniruzzaman*etal.* (2013), Tripathi*et al.* (2017),Jilani*et al.* (2010),Panwar*et al.* (2000) andAkoumianakis*et al*. (2011).

Table-1: Effect of nitrogen level on Plant height (cm),Number of leaves per plant and Fresh weight of leaves (gm)

Treatments	Plant height (cm)			Number of leaves per plant			Fresh weight of leaves (gm)		
	At 15 DAS	At 30 DAS	At Harvest	At 15 DAS	At 30 DAS	At Harvest	At 15 DAS	At 30 DAS	At Harvest
N_0- Control	7.20	16.38	35.30	4.38	6.16	8.96	26.25	37.76	60.21
N_1- 60 kg Nitrogen/ha	11.72	23.64	49.08	5.54	8.60	11.73	34.10	50.10	85.76
N_2- 80 kg Nitrogen/ha	12.66	25.12	53.48	5.70	8.85	12.76	38.68	64.55	97.55
N_3- 100 kg Nitrogen/ha	13.42	26.24	58.71	5.91	9.14	14.58	40.86	75.95	112.46
N_4- 120 kg Nitrogen/ha	13.84	26.50	65.10	6.72	9.84	17.10	44.10	83.28	118.32
N_5- 140 kg Nitrogen/ha	13.56	26.35	61.15	6.35	9.42	15.26	42.64	77.16	116.92
N_6- 160 kg Nitrogen/ha	12.96	25.45	56.07	5.82	8.98	13.47	40.12	69.21	105.83
N_7- 180 kg Nitrogen/ha	12.18	24.26	50.66	5.59	8.71	11.91	35.44	56.72	92.46
SEM(+/-)	**0.144**	**0.335**	**1.008**	**0.100**	**0.106**	**0.146**	**0.668**	**1.213**	**1.331**
C.D.at 5% of level	**0.436**	**1.016**	**3.058**	**0.302**	**0.323**	**0.444**	**2.025**	**3.678**	**4.036**

Table-2: Effect of nitrogen level on Leaf area per plant (cm^2) and Length of leaves at **harvesting (cm)**

Treatments	Leaf area per plant (cm^2)			Length of leaves at harvesting (cm)
	At 15 DAS	At 30 DAS	At Harvest	
N_0- Control	30.15	65.86	78.36	22.36
N_1- 60 kg Nitrogen/ha	36.95	88.52	97.10	29.10
N_2- 80 kg Nitrogen/ha	37.24	97.74	114.58	31.94
N_3- 100 kg Nitrogen/ha	39.78	120.38	155.12	35.45
N_4- 120 kg Nitrogen/ha	41.20	132.54	170.64	38.22
N_5- 140 kg Nitrogen/ha	40.36	128.12	167.33	36.58
N_6- 160 kg Nitrogen/ha	38.44	104.46	136.21	33.62
N_7- 180 kg Nitrogen/ha	37.10	94.75	105.29	30.54
SEM(+/-)	**0.588**	**1.495**	**2.041**	**0.504**
C.D.at 5% of level	**1.784**	**4.535**	**6.191**	**1.530**

Table-3: Effect of nitrogen level on Yield of Radish

Treatments	Root length (cm)	Root girth (cm)	Average plant weight at Harvesting (gm)	Fresh root weight (g/plant)	Yield per plot (kg)	Yield per hectare (q)
N_0- Control	13.94	5.21	124.42	105.14	15.26	168.24
N_1- 60 kg Nitrogen/ha	19.98	9.80	162.58	127.78	24.72	263.28
N_2- 80 kg Nitrogen/ha	21.54	10.48	184.72	138.92	27.56	304.85
N_3- 100 kg Nitrogen/ha	23.26	11.22	208.86	145.15	31.66	345.36
N_4- 120 kg Nitrogen/ha	26.78	12.10	224.15	150.45	34.15	376.52
N_5- 140 kg Nitrogen/ha	24.57	11.74	215.55	148.21	33.29	368.82
N_6- 160 kg Nitrogen/ha	22.45	10.86	192.21	143.35	29.48	324.58
N_7- 180 kg Nitrogen/ha	20.12	10.16	175.49	132.48	25.92	282.15
SEM(+/-)	**0.345**	**0.174**	**2.886**	**1.443**	**0.379**	**3.974**
C.D.at 5% of level	**1.045**	**0.529**	**8.752**	**4.376**	**1.149**	**12.053**

Conclusion:

Based on the above investigation, it can be concluded that there was a significant effect of nitrogen level on growth and yield attributing characters of radish cv. PusaChetki. Itisconducted that the treatment **N_4-** 120 kg Nitrogen/ha gave the highest growth and yield parameters *viz.*, Plant height (cm) at 15 DAS, 30 DAS and at harvest, numberofleavesperplantat 15 DAS, 30 DAS and at harvest, Fresh weight of leaves (gm) at 15 DAS, 30 DAS and at harvest, Leaf area per plant (cm^2) at 15 DAS, 30 DAS and at harvest, Length of leaves at harvesting (cm), Root length (cm), Root girth (cm), Average plant weight at Harvesting (gm), Fresh root weight (gm/plant), Yield per plot (kg) and Yield per hectare (q). So, we can suggest to farmers that the dose of 120 kg Nitrogen/ha gave the better results in term of growth and yield of Radish.

References:

1. Akoumianakis KA, Karapanos IC ,Giakoumaki M ,Alexopoulos AA and Passam HC.2011. Nitrogen, season and cultivar affect radish growth, yield, sponginess and hollowness , *International Journal of Plant Production,* 5(2):111-12[1]
2. Baloch PA,UddinR,NizamaniFK,Solangi AH and Siddiqui AA.2014.Effect of nitrogen, phosphorus and potassium fertilizers on growth and yield characteristics of radish *(Raphanussativus* L.),*American-Eurasian Journal of Agricultural & Environmental Sciences*, 14(6): 565-569.
3. Dash, S. K., Pathak, M., Tripathy, L. and Barik, S., 2018, Studies on effect of integrated nutrient management on growth and yield attributes in radish *(Raphanussativus*L.) and its residual effect in coriander (*CoriandrumSativum*L.) in radish-coriander cropping sequence *Pharmacognosy and Phytochemistry*, 8(1):319-322.
4. Hussain I, Haq I, Sajid M, *et al*. Effect of nitrogen alone and in combination with constant doses of potassium on yield of radish. Sarhad J Agric.1997;13:39–43.
5. JilaniMS ,Tariq B and Kashif W. 2010. Effect of nitrogen on growth and yield of radish, Journal of

Agricultural Research, 48(2): 219-225.

6. Kowalska I, Sady W, Shura A. Effect of of forms of nitrogen fertilizer, foliar feeding and place of cultivation on the yield and quality of lettuce. ActaAgrophysica. 2006;7(3):619–631.
7. KrishnkantN ,Sharma RK, Kushwah SS and Singh OP. 2018 . Influence of Varieties and Fertility Levels on Growth, Yield and Quality of Radish (Raphanussativus L.) Under Malwa Region of Madhya Pradesh , *International Journal of Agriculture Sciences* , 10(5) : 5371-5374.
8. MoniruzzamanM ,Akand MH, Hossain MI, Sarkar MD and Ullah A . 2013 .Effect of Nitrogen on the Growth and Yield of Carrot (*Daucuscarota* L.) , *A Scientific Journal of Krishi Foundation* ,11(1): 76-81.
9. PanwarAS ,Verma VS and Bawa R . 2000. Growth and seed yield of radish (Raphanussativus L.) as influenced by nitrogen and biofertilizer application, *Indian Journal of Agronomy*, 45 (2): 411-415.
10. Pathak M ,Tripathy P, Dash SK, Sahu GS and Pattanayak SK . 2017. Effect of source of nutrient on growth, yield and quality of Radish (*Raphanussativus* L.) in radish-coriander cropping sequence ,*The Pharma Innovation Journa*l , 6(12): 496-499.
11. Pervez MA, Ayub CM, Saleem BA. Effect of nitrogen levels and spacing on growth and yield of radish (*Raphanussativus*L.). Int J Agri Biol. 2004;06(3):504–506.
12. Poon TB, Regmi HN, Woli OB. Influence of plant spacing on seed yield of radish Mino Early. In: Khatri BB, Sharma BP, Khatiwada PP, *et al*, editors. Proceedings of the fourth national workshop on horticulture (2-4 March 2004). Khumaltar: Nepal Agri Res Council; 2004;373–376.
13. Sharma UG, Vihol NJ and ChavdaJC . 2013. Influence of plant density and nutrient management on growth, yield and quality of radish (*Raphanussativus*L.) cv. 'Pusachetki' ,The Asian Journal of Horticulture , 8(2) : 671-676.
14. Srinivas, K., &Naik, L. B. (1990). Growth and yield of radish (*Raphanussativus* L.) in relation to nitrogen and potash fertilization.

Indian Journal of Horticulture, 47(1), 114-119

15. SumagaysayCL .2014 . Comparative study of different levels of inorganic fertilizers on radish production, *International Journal of Educational Research* , 2(5): 185-194.
16. Thakar, MP, 2006, Effect of spacing and nitrogen on growth, yield and quality of radish *(Raphanussativus* L.), M.Sc. (Ag.) Thesis, JAU, Junagarh.
17. ThapaU ,Mohanto B , Chattopadhyay SB and Ghanti P. 2003. Growth and yield of some cultivars of radish (*Raphanussativus* L.) with nitrogen levels, *Environment and Ecology*, 21(4): 836-838
18. TripathiAK , Ram RB , Rout S, Kumar A and Patra SS . 2017 .Effect of Nitrogen Levels and Spacing on Growth and Yield of Radish *(Raphanussativus* L.) Cv. Kashi Sweta , *International Journal of Pure & Applied Bioscience*, 5(4):1951-1960 .
19. Yuan WL , Yuan S , Deng X , Gan C , Cui L and Wang Q . 2015 . Effect of N management on root yield and N uptake of radishes in southern china ,*Horticultural Science* , 50(5): 750-753.